

SARTINE ISLAND

ER #11

ORIGINAL PURPOSE To protect colonies of breeding seabirds and their habitat

OVERVIEW

Date established:	4 May 1971	Location:	33 km WNW of Cape Scott on the Cook Bank, between Beresford and Triangle Islands in the Scott Islands
ORC #:	3011		
Map number:	102 I/15		
Marine chart number:	3625		
		Latitude:	50°49'N
		Longitude:	128°54'W

Total Area:	1,091 ha	Elevation:	-60-105 m
Land:	30 ha		
Marine:	1,061 ha		

Access: Closed to public to protect nesting birds and their habitat. A permit is required to land on the island. It is accessible by boat, but landing is difficult.

Biogeoclimatic Zones:	Coastal Western Hemlock
Biogeoclimatic Variant:	CWFvh1 Southern Very Wet Hypermaritime
Terrestrial Ecoregion:	Vancouver Island Shelf
Marine Ecoregion:	Vancouver Island Shelf
Marine Ecoregions:	Benthic HHLFMCL (3 ha) Benthic HHLFMCM (1056 ha) Pelagic WE (569 ha)
Region:	Vancouver Island
Management Area:	Cape Scott

COMPOSITION

Physical: This is a rugged, elongate island which rises steeply from the sea to a series of pinnacles and humps. The shoreline is rock, rising directly from the water, except for a narrow boulder beach along the northeastern shore. Though steep, most of the upper slopes have a deep rich soil favourable for plant growth, and few vertical cliffs are present.

Biological: The island supports no trees, and shrubs are not extensive. Rather luxuriant stands of dune wildrye grass cover many slopes, and salmonberry is present at the summit where the ground is nearly flat in a few places. Interesting plant records include leatherly (formerly leatherleaf) polypody fern, hedge-nettle, Indian paintbrush, western buttercup and hairy rockcress. Twenty-three species of vascular plants have been collected or observed.

Nesting seabirds (766 000 birds, Canadian Wildlife Service, 1987) include Cassin's Auklet (376 000 pairs), Tufted Puffin (6400 pairs), Common Murre (135 pairs), Glaucous-winged Gull (240 pairs), Pelagic Cormorant (137 pairs) and Brandt's Cormorant (20 pairs). Additional cormorants, gulls and murre nest on rocks just outside the reserve. The Brandt's Cormorant colony, discovered in 1975, may in fact not be a breeding site, since this species is

known to build and attend “dummy” nests. Black Oystercatchers and Pigeon Guillemots (176 birds) nest around the shoreline. Fox and Song sparrows are the most common land birds. A pair of Peregrine Falcons nest here; Bald Eagles have been seen in summer and may nest as well. The only land mammal is a large, endemic sub-species of the Deer Mouse. Harbour Seals are common around the shoreline. Rocks just outside the reserve provide a summer rookery for 800-1200 Steller Sea Lions. Three species of isopods, one millipede and two land snails have been collected on Sartine Island. It is likely Sea Otter were common in the surrounding marine environment prior to extirpation. Occasional sightings are becoming more frequent indicating that Sea Otter are re-establishing colonies in this area.

MANAGEMENT CONCERNS

SIGNIFICANT SPECIES	BC LIST STATUS	COSEWIC STATUS	CF PRIORITY
Northern Abalone	Red listed	Threatened (2000)	2
Common Murre	Red listed		2
Brandt’s Cormorant	Red listed		1
Cassin’s Auklet	Blue listed		2
Tufted Puffin	Blue listed		2
Steller Sea Lion	Blue listed	Special Concern (2003)	2
Sea Otter	Blue listed	Special Concern (2007)	1
Peregrine Falcon spp. <i>pealei</i>	Blue listed	Special Concern (2007)	1

THREATS

Climate Change: The IPCC has projected an increase in sea surface temperature and raised sea levels, both of which could result in changes in the presence and life cycles of marine species which are important food sources for the protected seabird populations. The loss of synchrony between seabird phenology and seasonal food source availability has already been documented on Triangle Island as the probable cause of Cassin’s Auklet breeding failures.

Harvest: Commercial and recreational fishing disturbs marine ecosystems and shorebird activity. Vessel traffic in the vicinity can be the cause of mortality in seabirds and marine mammals especially as the source of chronic oiling.

Recreation: Close approach by marine or air traffic disrupts marine mammals and seabirds.

RESEARCH OPPORTUNITIES

Brief reports on seabird occurrence and abundance are available. Canadian Wildlife Service has surveyed vegetation cover. Environment Canada maintains a weather station on the island.
The marine environment has not been inventoried.
This site is ideal for continuing studies on climate change.

WARDEN ACTIVITIES

Monitor vegetation cover
Monitor intertidal communities

SCIENTIFIC NAMES OF SPECIES MENTIONED IN THE SARTINE ISLAND ER ACCOUNT**Flora**

buttercup, western (*Ranunculus occidentalis*)
fern, leatherly polypody (*Polypodium scoleri*)
hedge-nettle (*Stachys* spp)
paintbrush, Indian (*Castilleja* spp.)
rockcress, hairy (*Arabis hirsute*)
salmonberry (*Rubus spectabilis*)
wildrye, dune (*Elymus mollis*)

Fauna

Abalone Northern (*Haliotis kamtschatkana*)
Auklet, Cassin's (*Ptychamphus aleuticus*)
Cormorant, Brandt's (*Phalacrocorax penicillatus*)
Cormorant, Pelagic (*Phalacrocorax pelagicus*)
Deermouse, Sartine Island (*Peromyscus maniculatus sartinensis*)
Eagle, Bald (*Haliaeetus leucocephalus*)
Falcon, Peregrine (*Falco peregrinus*)
Guillemot, Pigeon (*Cephus columba*)
Gull, Glaucous-winged (*Larus glaucescens*)
Murre, Common (*Uria aalge*)
Oystercatcher, Black (*Haematopus bachmani*)
Puffin, Tufted (*Fratercula cirrhata*)
Sea Lion, Steller (*Eumetopias jubatus*)
Sea Otter (*Enhydra lutris*)
Seal Harbour (*Phoca vitulina*)
Sparrow, Fox (*Passerella iliaca*)
Sparrow, Song (*Melospiza melodia*)